



Our Lady Help of Christians Emergency response to Anaphylactic reaction



A First Aid Plan for Anaphylaxis poster is available on the [ASCIA website](#)

In all situations

1. If safe to do so, lay the person flat, do not allow patient to stand or walk.
2. If breathing is difficult allow patient to sit
 - Be calm, reassuring
 - Do not leave them alone.
 - Seek assistance from another staff member or reliable student to locate the autoinjector or a general use autoinjector, and the student's Individual Anaphylaxis Management Plan
 - If the student appears to be experiencing a first time reaction, continue with steps 2 – 6.
3. Administer prescribed adrenaline autoinjector – note the time given and retain used EpiPen to give ambulance paramedics.
4. Phone ambulance 000 (112 – mobile).
5. If there is no improvement or severe symptoms progress, further adrenaline doses may be given every five minutes (if another autoinjector is available).
6. Phone family/emergency contact.

If in doubt, give an autoinjector

If the student has not been previously diagnosed with an allergy or at risk of anaphylaxis but appears to be having a severe allergic reaction phone ambulance 000 (112 – mobile), advise them that the school has a general use autoinjector and seek and follow the advice given by them.

Approval authority	Director, Learning and Regional Services
Approval date	5 July 2023
Next review	March 2025